

Diophantine Approximations and the Convergence of Certain Series

Alexander Begunts, Dmitry Goryashin

Abstract

Consider two series

$$\sum_{n=1}^{\infty} \frac{\sin^n \pi \theta n}{n^\alpha}, \quad \sum_{n=1}^{\infty} \frac{\cos^n \pi \theta n}{n^\alpha}.$$

We show that number-theoretical properties of θ have a strong effect on the convergence when $0 < \alpha \leq 1$. The complete investigation for $\theta \in \mathbb{Q}$ is given. For irrational θ we prove the result which depends on how well θ can be approximated with rational numbers, i.e. on its irrationality measure. We obtain that if $\alpha > \frac{1}{2}$ then both series converge absolutely for almost all real θ . Finally, we construct such an everywhere dense set of θ that both series diverge when $\alpha \leq 1$.

Introduction

Various authors considered series which converge or diverge depending on number-theoretical properties of their parameters. A century ago G.H. Hardy and J.E. Littlewood [4], *inter alia*, investigated the series

$$\sum_{n=1}^{\infty} \frac{\sin \pi \theta n^2}{n^\alpha}, \quad \sum_{n=1}^{\infty} \frac{\cos \pi \theta n^2}{n^\alpha}.$$

It follows from their results that these series diverge for all irrational θ when $\alpha \leq \frac{1}{2}$, and converge for all irrational numbers θ with bounded partial quotients (for example, quadratic irrationals) when $\alpha > \frac{1}{2}$.¹ They also constructed examples of irrational θ , for which both series diverge for all $\alpha \leq 1$.

G.I. Arkhipov and K.I. Oskolkov [1] considered the series

$$\sum_{n=1}^{\infty} \frac{\sin \pi \theta n^k}{n}.$$

Their brilliant result states that it converges *for all* real θ if k is odd. Moreover, they point out such values of θ for which the series diverges for any even k .

In 2011, E. Laeng and V. Pata [7] presented a convergence-divergence test for series of non-negative terms which fail to be monotonic. As an application of the test the authors prove the convergence of one series involving the linear function of $\cos n$ raised to the n -th power, using the fact that the irrationality measure of π is finite.

In this paper we combine methods of analysis and number theory to investigate two series

$$\sum_{n=1}^{\infty} \frac{\sin^n \pi \theta n}{n^\alpha}, \quad \sum_{n=1}^{\infty} \frac{\cos^n \pi \theta n}{n^\alpha}.$$

¹In fact, it follows from estimates of H. Weyl sums, obtained later, and celebrated Roth's theorem, that both series converge for all irrational algebraic θ when $\alpha > \frac{1}{2}$.

Since they converge absolutely for $\alpha > 1$, from now on we keep in view the case $\alpha \leq 1$. It will be shown that number-theoretical properties of θ have a strong effect on the convergence. In case of rational θ we prove the following statements.

Theorem 1. *The series*

$$\sum_{n=1}^{\infty} \frac{\cos^n \pi \theta n}{n^\alpha},$$

where $\theta = \frac{p}{q}$ with coprime integers p, q , and $0 < \alpha \leq 1$,

- 1) diverges to $+\infty$ if at least one of p, q is even;
- 2) converges conditionally if both p, q are odd.

Theorem 2. *The series*

$$\sum_{n=1}^{\infty} \frac{\sin^n \pi \theta n}{n^\alpha},$$

where $\theta = \frac{p}{q}$ with coprime integers p, q , and $0 < \alpha \leq 1$,

- 1) converges absolutely if q is odd;
- 2) converges conditionally if q is even and not divisible by 4;
- 3) diverges to $+\infty$ if q is divisible by 4.

Next, we proceed to the case of irrational θ . Recall the following definition from the theory of Diophantine approximations: the irrationality measure (or irrationality exponent) of real number θ is the infimum $\mu(\theta)$ of the set of positive real numbers μ for which

$$0 < \left| \theta - \frac{p}{q} \right| < \frac{1}{q^\mu}$$

holds for (at most) finitely many solutions p/q with p and q integers. If no such μ exists we set $\mu(\theta) = +\infty$ and say that θ is a Liouville number.

It follows from this definition that if θ is a number with finite irrationality measure μ , then for all $\delta > 0$ there exists a constant $c = c(\theta, \delta) > 0$ such that the inequality

$$\left| \theta - \frac{p}{q} \right| > \frac{c}{q^{\mu+\delta}}$$

is true for all integers p and $q \geq 1$.

Note that $\mu(\theta) = 1$ if and only if θ is rational, and famous Roth's theorem [2, Chapter VI] states that for irrational algebraic numbers it holds $\mu(\theta) = 2$. For transcendental θ we have $\mu(\theta) \geq 2$.

Theorem 3. *Let θ be an irrational number with finite irrationality measure $\mu = \mu(\theta)$. If $\alpha > 1 - \frac{1}{2\mu-2}$, then both series*

$$\sum_{n=1}^{\infty} \frac{\sin^n \pi \theta n}{n^\alpha}, \quad \sum_{n=1}^{\infty} \frac{\cos^n \pi \theta n}{n^\alpha}$$

converge absolutely.

As a corollary we immediately obtain that both series converge absolutely for any irrational algebraic number θ when $\alpha > \frac{1}{2}$. For instance, using the well known result $\mu(e) = 2$ and the recently obtained estimate $\mu(1/\pi) = \mu(\pi) \leq 7.6063 \dots$ from [8] we see that both series

$$\sum_{n=1}^{\infty} \frac{\sin^n(\pi e n)}{n^\alpha}, \quad \sum_{n=1}^{\infty} \frac{\cos^n(\pi e n)}{n^\alpha}$$

converge absolutely for $\alpha > \frac{1}{2}$, and both series

$$\sum_{n=1}^{\infty} \frac{\sin^n n}{n^\alpha}, \quad \sum_{n=1}^{\infty} \frac{\cos^n n}{n^\alpha}$$

converge absolutely for $\alpha > 0.9243 \dots$

Moreover, the key result of continued fractions metric theory (see [5, § 14, Theorem 32]) states that $\mu(\theta) = 2$ holds for almost all real θ . Therefore from Theorem 3 one can deduce the upcoming assertion.

Theorem 4. *Let $\alpha > \frac{1}{2}$. Then both series*

$$\sum_{n=1}^{\infty} \frac{\sin^n \pi \theta n}{n^\alpha}, \quad \sum_{n=1}^{\infty} \frac{\cos^n \pi \theta n}{n^\alpha}$$

converge absolutely for almost all real θ .

Yet, we submit another (and more simple) proof which is independent of Theorem 3 and involves methods of real analysis.

In addition to the investigation of the case of irrational θ , we construct the set of transcendental numbers for which both series converge only for $\alpha > 1$.

Theorem 5. *There exists such an everywhere dense in $\mathbb{R}$ set of irrational numbers E , that for any $\xi \in E$ both series*

$$\sum_{n=1}^{\infty} \frac{\sin^n \pi \xi n}{n^\alpha}, \quad \sum_{n=1}^{\infty} \frac{\cos^n \pi \xi n}{n^\alpha}$$

diverge for all $\alpha \leq 1$.

1 Proof of Theorem 1

1) Decompose the series into the sum of q series each corresponding to values $n \equiv a \pmod{q}$, $a = 1, 2, \dots, q$:

$$\sum_{n=1}^{\infty} \frac{\cos^n \frac{\pi p n}{q}}{n^\alpha} = \sum_{a=1}^q \sum_{l=0}^{\infty} \frac{\left(\cos \frac{\pi p(lq+a)}{q} \right)^{lq+a}}{(lq+a)^\alpha} = \sum_{a=1}^q \sum_{l=0}^{\infty} \frac{\left(\cos \left(\frac{\pi a p}{q} + \pi p l \right) \right)^{lq+a}}{(lq+a)^\alpha}.$$

We start with p even, $p = 2p_1$. Since $(p, q) = 1$, we obtain that q is odd. Among the numbers $\cos \frac{2\pi a p_1}{q}$, $a = 1, 2, \dots, q$, all but one lie in the interval $(-1; 1)$. Indeed, if $a = q$, then $\cos \frac{2\pi a p_1}{q} = \cos 2\pi p_1 = 1$; for $1 \leq a < q$ we have $\frac{2\pi a p_1}{q}$ noninteger, hence $\cos \frac{2\pi a p_1}{q} \neq \pm 1$. Thus in this case the series can be written as

$$\sum_{a=1}^q \sum_{l=0}^{\infty} \frac{\left(\cos \frac{2\pi a p_1}{q} \right)^{lq+a}}{(lq+a)^\alpha} = \sum_{a=1}^{q-1} \cos^a \frac{2\pi a p_1}{q} \sum_{l=0}^{\infty} \frac{\left(\cos^q \frac{2\pi a p_1}{q} \right)^l}{(lq+a)^\alpha} + \sum_{l=0}^{\infty} \frac{1}{q^\alpha (l+1)^\alpha},$$

Since $|\cos^q \frac{2\pi a p_1}{q}| < 1$, the first $q-1$ series in the sum converge absolutely, and the last series diverges for $0 < \alpha \leq 1$. Therefore the series under consideration diverges to $+\infty$.

Next, let $q = 2q_1$ and p is odd. The equality $\cos(\frac{\pi a p}{2q_1} + \pi p l) = \pm 1$ is equivalent to $ap \equiv 0 \pmod{2q_1}$ which holds only when $a = q = 2q_1$. Hence the series can also be represented as

$$\sum_{a=1}^{2q_1} \sum_{l=0}^{\infty} \frac{\left(\cos \left(\frac{\pi a p}{2q_1} + \pi p l \right) \right)^{2lq_1+a}}{(2lq_1+a)^\alpha} =$$

$$= \sum_{a=1}^{2q_1-1} \sum_{l=0}^{\infty} \frac{\left(\cos^{2q_1} \frac{\pi ap}{2q_1}\right)^l \cos^a\left(\frac{\pi ap}{2q_1} + \pi pl\right)}{(2lq_1 + a)^\alpha} + \sum_{l=0}^{\infty} \frac{1}{q^\alpha(l+1)^\alpha}.$$

In the same way as in the previous case we observe that the series diverges to $+\infty$.

2) Let both p, q be odd. Decompose the series into the sum of $2q$ series each corresponding to values $n \equiv a \pmod{2q}$, $a = 1, 2, \dots, 2q$:

$$\sum_{n=1}^{\infty} \frac{\cos^n \frac{\pi pn}{q}}{n^\alpha} = \sum_{a=1}^{2q} \sum_{l=0}^{\infty} \frac{\left(\cos \frac{\pi p(2lq+a)}{q}\right)^{2lq+a}}{(2lq+a)^\alpha} = \sum_{a=1}^{2q} \sum_{l=0}^{\infty} \frac{\left(\cos \frac{\pi ap}{q}\right)^{2lq+a}}{(2lq+a)^\alpha}.$$

The equality $\cos \frac{\pi ap}{q} = 1$ is equivalent to $ap \equiv 0 \pmod{2q}$ which holds only when $a = 2q$, because p and $2q$ are coprime. Next, $\cos \frac{\pi ap}{q} = -1$ means that $ap \equiv q \pmod{2q}$, which has a unique solution $a = a_0$, $1 \leq a_0 \leq 2q-1$. Note that a_0 is odd. Hence $|\cos \frac{\pi ap}{q}| < 1$ whenever $a \neq a_0$ and $a \neq 2q$. Rewriting the series as

$$\begin{aligned} & \sum_{a=1}^{2q} \sum_{l=0}^{\infty} \frac{\left(\cos \frac{\pi ap}{q}\right)^{2lq+a}}{(2lq+a)^\alpha} = \\ &= \sum_{\substack{a=1 \\ a \neq a_0}}^{2q-1} \cos^a \frac{\pi ap}{q} \sum_{l=0}^{\infty} \frac{\left(\cos^{2q} \frac{\pi ap}{q}\right)^l}{(2lq+a)^\alpha} + \sum_{l=0}^{\infty} \left(\frac{1}{(2ql+2q)^\alpha} - \frac{1}{(2ql+a_0)^\alpha} \right), \end{aligned}$$

we conclude that $2q-2$ series in the first sum converge absolutely. Now

$$\begin{aligned} & \frac{1}{(2ql+2q)^\alpha} - \frac{1}{(2ql+a_0)^\alpha} = \frac{1}{(2ql)^\alpha} \left(\left(1 + \frac{1}{l}\right)^{-\alpha} - \left(1 + \frac{a_0}{2ql}\right)^{-\alpha} \right) = \\ &= \frac{1}{(2ql)^\alpha} \left(1 - \frac{\alpha}{l} + O\left(\frac{1}{l}\right) - 1 + \frac{\alpha a_0}{2ql} + O\left(\frac{1}{l}\right) \right) \sim \frac{\alpha\left(\frac{a_0}{2q} - 1\right)}{(2q)^\alpha} \cdot \frac{1}{l^{\alpha+1}} \end{aligned}$$

for $l \rightarrow \infty$. Since $\alpha + 1 > 1$, the series converges. It remains to be seen that the series of absolute values

$$\begin{aligned} & \sum_{a=1}^{2q} \sum_{l=0}^{\infty} \frac{\left|\cos \frac{\pi ap}{q}\right|^{2lq+a}}{(2lq+a)^\alpha} = \\ &= \sum_{\substack{a=1 \\ a \neq a_0}}^{2q-1} \cos^a \frac{\pi ap}{q} \sum_{l=0}^{\infty} \frac{\left(\cos^{2q} \frac{\pi ap}{q}\right)^l}{(2lq+a)^\alpha} + \sum_{l=0}^{\infty} \left(\frac{1}{(2ql+2q)^\alpha} + \frac{1}{(2ql+a_0)^\alpha} \right), \end{aligned}$$

evidently diverges, thus the series under consideration converges conditionally.

2 Proof of Theorem 2

By analogy with the previous proof, decompose the series into the sum of $2q$ series:

$$\sum_{n=1}^{\infty} \frac{\sin^n \frac{\pi pn}{q}}{n^\alpha} = \sum_{a=1}^{2q} \sum_{l=0}^{\infty} \frac{\left(\sin \frac{\pi p(2lq+a)}{q}\right)^{2lq+a}}{(2lq+a)^\alpha} = \sum_{a=1}^{2q} \sum_{l=0}^{\infty} \frac{\left(\sin \frac{\pi ap}{q}\right)^{2lq+a}}{(2lq+a)^\alpha}.$$

1) If q is odd, then the equality $\frac{\pi ap}{q} = \frac{\pi}{2} + \pi k$ means that $2ap = q + 2kq$, which is impossible for any integer k . Hence in this case $|\sin \frac{\pi ap}{q}| < 1$ for all $a = 1, 2, \dots, 2q$, thus all the series in the right-hand side converge absolutely, and so does the series under study.

2) Let $q = 2q_1$, where q_1 is odd. Then the equality $\frac{\pi ap}{2q_1} = \frac{\pi}{2} + 2\pi k$ holds if and only if $ap \equiv q_1 \pmod{4q_1}$. Since $(p, 4q_1) = 1$, the congruence has a unique solution $a \equiv a_0 \pmod{4q_1}$ where $1 \leq a_0 < 4q_1 = 2q$. In a similar way, $\frac{\pi ap}{2q_1} = -\frac{\pi}{2} + 2\pi k$ is equivalent to $ap \equiv -q_1 \pmod{4q_1}$, which is true only for $a = 2q - a_0$. Note that a_0 is odd. Then for $a \neq a_0, a \neq 2q - a_0$ it holds $|\sin \frac{\pi ap}{q}| < 1$, so we can rewrite the series as

$$\sum_{\substack{a=1 \\ a \neq a_0 \\ a \neq 2q-a_0}}^{2q} \sum_{l=0}^{\infty} \frac{\left(\sin \frac{\pi ap}{q}\right)^{2lq+a}}{(2lq+a)^\alpha} + \sum_{l=0}^{\infty} \left(\frac{1}{(2lq+a_0)^\alpha} - \frac{1}{(2lq+2q-a_0)^\alpha} \right).$$

and see that $2q-2$ series in the first sum converge absolutely. The last series is also convergent as far as

$$\begin{aligned} \frac{1}{(2ql+a_0)^\alpha} - \frac{1}{(2ql+2q-a_0)^\alpha} &= \frac{1}{(2ql)^\alpha} \left(\left(1 + \frac{a_0}{2ql}\right)^{-\alpha} - \right. \\ &\left. - \left(1 + \frac{2q-a_0}{2ql}\right)^{-\alpha} \right) = \frac{1}{(2ql)^\alpha} \left(1 - \frac{a_0\alpha}{2ql} + O\left(\frac{1}{l}\right) - 1 + \frac{\alpha(2q-a_0)}{2ql} + \right. \\ &\left. + O\left(\frac{1}{l}\right) \right) \sim \frac{\alpha(1 - \frac{a_0}{q})}{(2q)^\alpha} \cdot \frac{1}{l^{\alpha+1}} \quad \text{when } l \rightarrow \infty. \end{aligned}$$

It converges conditionally, since the series of absolute values

$$\begin{aligned} &\sum_{n=1}^{\infty} \frac{|\sin \frac{\pi pn}{q}|^n}{n^\alpha} = \\ &= \sum_{\substack{a=1 \\ a \neq a_0 \\ a \neq 2q-a_0}}^{2q} \sum_{l=0}^{\infty} \frac{\left|\sin \frac{\pi ap}{q}\right|^{2lq+a}}{(2lq+a)^\alpha} + \sum_{l=0}^{\infty} \left(\frac{1}{(2lq+a_0)^\alpha} + \frac{1}{(2lq+2q-a_0)^\alpha} \right) \end{aligned}$$

diverges.

3) After all, let $q = 4q_1$. Then the equality $\frac{\pi ap}{4q_1} = \frac{\pi}{2} + \pi k$ is equivalent to $ap \equiv 2q_1 \pmod{4q_1}$. Since $(p, 4q_1) = 1$, the congruence has a unique solution $a \equiv a_0 \pmod{4q_1}$ where $1 \leq a_0 < 4q_1 = q$ and a_0 is even. Thus the series under study can be rewritten as

$$\sum_{\substack{a=1 \\ a \neq a_0 \\ a \neq q+a_0}}^{2q} \sum_{l=0}^{\infty} \frac{\left(\sin \frac{\pi ap}{q}\right)^{2lq+a}}{(2lq+a)^\alpha} + \sum_{l=0}^{\infty} \left(\frac{1}{(2lq+a_0)^\alpha} + \frac{1}{(2lq+q+a_0)^\alpha} \right),$$

and therefore diverges to $+\infty$. This completes the proof.

3 Proof of Theorem 3

We assume that $\theta > 0$ and prove that the series

$$\sum_{n=1}^{\infty} \frac{|\cos(\pi n\theta)|^n}{n^\alpha}$$

converges for $\alpha > 1 - \frac{1}{2\mu-2}$. The sin-series can be considered in the same way. Since this is the series of absolute values, by the well-known fact its convergence is independent of the permutation of its terms. Thus the series can be represented as

$$\sum_{n=1}^{\infty} \frac{|\cos(\pi n \theta)|^n}{n^\alpha} = \sum_{s=0}^{\infty} \sum_{\substack{n \in \mathbb{N} \\ 1 - \frac{1}{2^s} \leq |\cos(\pi n \theta)| < 1 - \frac{1}{2^{s+1}}}} \frac{|\cos(\pi n \theta)|^n}{n^\alpha} = \sum_{s=0}^{\infty} S_{\frac{1}{2^s}},$$

where

$$S_\varepsilon = \sum_{\substack{n \in \mathbb{N} \\ 1 - \varepsilon \leq |\cos(\pi n \theta)| < 1 - \varepsilon/2}} \frac{|\cos(\pi n \theta)|^n}{n^\alpha}.$$

We need an upper estimate for S_ε , where ε is fixed and small enough (greater values of ε have an effect on at most finitely many terms of the series $\sum_{s=0}^{\infty} S_{\frac{1}{2^s}}$). For that purpose we prove the following statement.

Lemma 3.1. *Let θ be a positive irrational number with finite irrationality measure $\mu = \mu(\theta)$, $\nu = \frac{1}{2\mu-2}$, $0 < \varepsilon < \min(1; 1 - \cos \frac{\pi\theta}{2})$, and let $\{n_k\}_{k=1}^{\infty}$ be the increasing sequence of positive integers for which*

$$1 - \varepsilon \leq |\cos(\pi n \theta)| < 1 - \varepsilon/2.$$

Then for all $\delta \in (0; \nu)$ there exists such a number $A = A(\theta, \delta) > 0$ (independent of ε) that for all positive integers k the following inequality holds:

$$n_k \geq Ak\varepsilon^{-\nu+\delta}.$$

Proof. Let $m = n_k$ and $n = n_{k+1}$ be two neighbor terms of $\{n_k\}_{k=1}^{\infty}$. Since $|\cos(\pi n \theta)| \geq 1 - \varepsilon$ and $|\cos(\pi m \theta)| \geq 1 - \varepsilon$, we obtain

$$\begin{aligned} -\arccos(1 - \varepsilon) + \pi l_n &\leq \pi n \theta \leq \arccos(1 - \varepsilon) + \pi l_n, \\ -\arccos(1 - \varepsilon) + \pi l_m &\leq \pi m \theta \leq \arccos(1 - \varepsilon) + \pi l_m \end{aligned}$$

for some $l_n, l_m \in \mathbb{N}$, so

$$|n\theta - l_n| \leq \frac{1}{\pi} \arccos(1 - \varepsilon), \quad |m\theta - l_m| \leq \frac{1}{\pi} \arccos(1 - \varepsilon).$$

Then

$$|(n - m)\theta - l| \leq |n\theta - l_n| + |m\theta - l_m| \leq \frac{2}{\pi} \arccos(1 - \varepsilon),$$

where $l = l_n - l_m \geq 0$. Note that if $\varepsilon < \min(1; 1 - \cos \frac{\pi\theta}{2})$ then $l_n = l_m$ can hold only when $(n - m)\theta < \frac{2}{\pi} \arccos(1 - \varepsilon) < \frac{2}{\pi} \frac{\pi\theta}{2} = \theta$, i.e. $n - m < 1$, a contradiction. Hence, $l = l_n - l_m > 0$. Dividing by $l\theta > 0$ we get an estimate

$$\left| \frac{n - m}{l} - \frac{1}{\theta} \right| \leq \frac{2}{l\pi\theta} \arccos(1 - \varepsilon) < \frac{4\sqrt{\varepsilon}}{l\pi\theta}.$$

Here we also used the inequality $\arccos(1 - \varepsilon) < 2\sqrt{\varepsilon}$ which holds when $0 < \varepsilon < 1$. Indeed, for the function $f(x) = \arccos(1 - x) - 2\sqrt{x}$ we have $f'(x) = \frac{1}{\sqrt{x(2-x)}} - \frac{1}{\sqrt{x}} < 0$ when $x \in (0; 1)$, so $f(x)$ is decreasing in this interval, thus $f(x) < f(0) = 0$.

Alternately, $\frac{1}{\theta}$ has a finite irrationality measure μ , thus for all $\delta' > 0$ the inequality

$$\left| \frac{n - m}{l} - \frac{1}{\theta} \right| > \frac{c}{l^{\mu+\delta'}}$$

is true for all integers $l \geq 1$, where $c = c(\theta, \delta')$. Hence we have

$$\frac{c}{l^{\mu+\delta'}} < \frac{4\sqrt{\varepsilon}}{l\pi\theta}, \quad l > \left(\frac{\pi\theta c}{4\sqrt{\varepsilon}} \right)^{\frac{1}{\mu-1+\delta'}}.$$

Therefore,

$$n_{k+1} - n_k = n - m \geq \frac{l-1}{\theta} \geq \frac{l}{2\theta} > A\varepsilon^{-\frac{1}{2\mu-2+2\delta'}} = A\varepsilon^{-\nu+\delta},$$

where $\delta' = \frac{\delta}{2\nu(\nu-\delta)} > 0$ and $A = \frac{1}{2\theta} \left(\frac{\pi\theta c}{4} \right)^{\frac{1}{\mu-1+\delta'}}$ is independent of ε .

Finally, we note that an analogous argument leads to the same estimate for n_1 . Thus the statement of lemma follows by induction. $\square$

Next, we apply Lemma 3.1 to handle S_ε . Since $n_k \geq Ak\varepsilon^{-\nu}$ we have

$$\begin{aligned} S_\varepsilon &= \sum_{\substack{n \in \mathbb{N} \\ 1-\varepsilon \leq |\cos(\pi n \theta)| < 1-\varepsilon/2}} \frac{|\cos(\pi n \theta)|^n}{n^\alpha} = \sum_{k=1}^{\infty} \frac{|\cos(\pi n_k \theta)|^{n_k}}{n_k^\alpha} \ll \\ &\ll \sum_{k=1}^{\infty} \frac{(1 - \frac{\varepsilon}{2})^{n_k}}{n_k^\alpha} \leq \sum_{k=1}^{\infty} \frac{(1 - \frac{\varepsilon}{2})^{Ak\varepsilon^{-\nu+\delta}}}{(Ak\varepsilon^{-\nu+\delta})^\alpha} = \frac{1}{A^\alpha \varepsilon^{-(\nu+\delta)\alpha}} \sum_{k=1}^{\infty} \frac{\left((1 - \frac{\varepsilon}{2})^{A\varepsilon^{-\nu+\delta}} \right)^k}{k^\alpha}. \end{aligned}$$

For $\alpha = 1$ and $\alpha < 1$ we suggest different approaches.

Let $\alpha = 1$. Since

$$\sum_{k=1}^{\infty} \frac{z^k}{k} = -\ln(1-z), \quad |z| < 1,$$

we set $z = (1 - \frac{\varepsilon}{2})^{A\varepsilon^{-\nu+\delta}}$ and obtain

$$\begin{aligned} S_\varepsilon &\ll -\frac{1}{A\varepsilon^{-\nu+\delta}} \ln \left(1 - (1 - \frac{\varepsilon}{2})^{A\varepsilon^{-\nu+\delta}} \right) = -\frac{1}{A\varepsilon^{-\nu+\delta}} \ln \left(1 - e^{A\varepsilon^{-\nu+\delta} \ln(1-\frac{\varepsilon}{2})} \right) = \\ &= -\frac{1}{A\varepsilon^{-\nu+\delta}} \ln \left(1 - e^{-\frac{A}{2}\varepsilon^{-\nu+\delta+1}(1+o(1))} \right) = -\frac{1}{A\varepsilon^{-\nu+\delta}} \ln \left(\frac{A}{2}\varepsilon^{-\nu+\delta+1}(1+o(1)) \right) \sim \\ &\sim -\frac{(1-\nu+\delta)\ln\varepsilon}{A\varepsilon^{-\nu+\delta}} \quad \text{when } \varepsilon \rightarrow +0. \end{aligned}$$

Hence the series

$$\sum_{n=1}^{\infty} \frac{|\cos(\pi n \theta)|^n}{n} = \sum_{s=0}^{\infty} S_{\frac{1}{2^s}} \ll \sum_{s=0}^{\infty} \frac{(1-\nu+\delta)s \ln 2}{A(2^{\nu-\delta})^s}$$

is convergent because $2^{\nu-\delta} = 2^{\frac{1}{2\mu-2}-\delta} > 1$ for sufficiently small $\delta > 0$.

Now let $\alpha < 1$. We formulate a special case of an asymptotic formula from [3, § 4, paragraph 3] as a lemma.

Lemma 3.2. *For $\alpha < 1$ it holds*

$$\sum_{k=1}^{\infty} \frac{z^k}{k^\alpha} \sim \Gamma(1-\alpha) \left(\ln \frac{1}{z} \right)^{\alpha-1} \quad \text{when } z \rightarrow 1-0.$$

Again put $z = (1 - \frac{\varepsilon}{2})^{A\varepsilon^{-\nu+\delta}}$ and for $\alpha < 1$ we have

$$\begin{aligned} S_\varepsilon &\ll \frac{1}{A^\alpha \varepsilon^{(-\nu+\delta)\alpha}} \Gamma(1-\alpha) (-A\varepsilon^{-\nu+\delta} \ln(1 - \frac{\varepsilon}{2}))^{\alpha-1} \sim \\ &\sim \frac{\Gamma(1-\alpha)}{2^{\alpha-1} A} \varepsilon^{(1-\nu+\delta)(\alpha-1)-(-\nu+\delta)\alpha} = \frac{\Gamma(1-\alpha)}{2^{\alpha-1} A} \varepsilon^{\alpha+\nu-\delta-1} \quad \text{when } \varepsilon \rightarrow +0. \end{aligned}$$

Thus the series

$$\sum_{n=1}^{\infty} \frac{|\cos(\pi n \theta)|^n}{n^\alpha} = \sum_{s=1}^{\infty} S_{\frac{1}{2^s}} \ll \frac{\Gamma(1-\alpha)}{2^{\alpha-1} A} \sum_{s=1}^{\infty} \frac{1}{(2^{\alpha+\nu-\delta-1})^s}$$

converges if $2^{\alpha+\nu-\delta-1} > 1$, i.e. if $\alpha > 1 - \nu + \delta = 1 - \frac{1}{2\mu-2} + \delta$ for sufficiently small $\delta > 0$. Proof of Theorem 3 is completed.

4 Analytic proof of Theorem 4

For the reason of periodicity it is enough to prove the convergence for almost all real θ from $[0; 1]$ when $\alpha > \frac{1}{2}$. We consider the sin-series; one can easily handle the cos-series in a similar way. Since

$$\int_0^1 \frac{|\sin(\pi n \theta)|^n}{n^\alpha} d\theta = \frac{1}{\pi n^{\alpha+1}} \int_0^{\pi n} |\sin x|^n dx = \frac{2n}{\pi n^{\alpha+1}} \int_0^{\pi/2} \sin^n x dx = \frac{2I_n}{\pi n^\alpha},$$

where

$$I_n = \int_0^{\pi/2} \sin^n x dx \asymp \frac{1}{\sqrt{n}},$$

it holds

$$\int_0^1 \frac{|\sin(\pi n \theta)|^n}{n^\alpha} d\theta \asymp \frac{1}{n^{\alpha+\frac{1}{2}}},$$

hence the series with integral terms

$$\sum_{n=1}^{\infty} \int_0^1 \frac{|\sin(\pi n \theta)|^n}{n^\alpha} d\theta$$

converges for $\alpha + \frac{1}{2} > 1$, i.e. for $\alpha > \frac{1}{2}$. By the corollary of B. Levi's monotone convergence theorem [6, sec. 30.1] it follows that the series

$$\sum_{n=1}^{\infty} \frac{|\sin(\pi n \theta)|^n}{n^\alpha}$$

converges for $\alpha > \frac{1}{2}$ almost everywhere on $[0; 1]$ and therefore on $\mathbb{R}$.

5 Divergence of the series on an everywhere dense set

In this section we reveal that every interval contains irrational numbers θ with infinite irrationality measure for which both series diverge for all $\alpha \leq 1$. This shows that the result analogous to Theorem 3 doesn't hold for such numbers. First, we need the following lemma on the rate of divergence of the series under consideration for some rational θ .

Lemma 5.1. Let $\theta = \frac{p}{q}$ with coprime numbers p, q , where q is divisible by 4. Then there exists such a positive integer A_q that the following estimates hold uniformly for all positive integers L and $\alpha \leq 1$:

$$\left| \sum_{n=1}^{2qL} \frac{\sin^n \frac{\pi p n}{q}}{n^\alpha} \right| \geq \frac{1}{q} \ln L - A_q, \quad \left| \sum_{n=1}^{2qL} \frac{\cos^n \frac{\pi p n}{q}}{n^\alpha} \right| \geq \frac{1}{q} \ln L - A_q,$$

Proof. In the proof of Theorem 2 we established the following representation of the sin-series partial sum:

$$\sum_{n=1}^{N_1} \frac{\sin^n \frac{\pi p n}{q}}{n^\alpha} = \sum_{\substack{a=1 \\ a \neq a_0 \\ a \neq q+a_0}}^{2q} \sum_{l=0}^L \frac{\left(\sin \frac{\pi a p}{q} \right)^{2lq+a}}{(2lq+a)^\alpha} + \sum_{l=0}^L \left(\frac{1}{(2lq+a_0)^\alpha} + \frac{1}{(2lq+q+a_0)^\alpha} \right).$$

Since for $a \neq a_0$, $a \neq q+a_0$ all the series

$$\sum_{l=0}^{\infty} \frac{\left(\sin \frac{\pi a p}{q} \right)^{2lq+a}}{(2lq+a)^\alpha}$$

converge absolutely for all α , there exists such an integer A_q depending only on q that

$$\left| \sum_{\substack{a=1 \\ a \neq a_0 \\ a \neq q+a_0}}^{2q} \sum_{l=0}^L \frac{\left(\sin \frac{\pi a p}{q} \right)^{2lq+a}}{(2lq+a)^\alpha} \right| \leq \sum_{\substack{a=1 \\ a \neq a_0 \\ a \neq q+a_0}}^{2q} \sum_{l=0}^{\infty} \frac{\left| \sin \frac{\pi a p}{q} \right|^{2lq+a}}{(2lq+a)^\alpha} \leq A_q.$$

Now it can be easily seen that

$$\begin{aligned} \sum_{l=0}^{\infty} \left(\frac{1}{(2lq+a_0)^\alpha} + \frac{1}{(2lq+q+a_0)^\alpha} \right) &\geq \sum_{l=0}^L \left(\frac{1}{2lq+2q} + \frac{1}{2lq+q+q} \right) = \\ &= \frac{1}{q} \sum_{l=0}^L \frac{1}{l+1} \geq \frac{1}{q} \sum_{l=1}^L \frac{1}{l} \geq \frac{1}{q} \ln L. \end{aligned}$$

Thus

$$\left| \sum_{n=1}^{2qL} \frac{\sin^n \frac{\pi p n}{q}}{n^\alpha} \right| \geq \frac{1}{q} \ln L - A_q.$$

The analogous inequality for the cos-series can be proved in the same way using proof of Theorem 1. $\square$

Proof of Theorem 5. Our goal is to show that every interval $(x_1; x_2) \subset \mathbb{R}$ contains such an irrational number ξ that both series

$$\sum_{n=1}^{\infty} \frac{\sin^n \pi \xi n}{n^\alpha}, \quad \sum_{n=1}^{\infty} \frac{\cos^n \pi \xi n}{n^\alpha}$$

diverge for all $\alpha \leq 1$. This construction uses the idea similar to the one implemented in [4]. We start with choosing an integer u and positive integers $b_1 < b_2 < \dots < b_s < \nu_1$ for which both numbers

$$\xi_0 = u + \sum_{i=1}^s \frac{1}{2^{b_i}} \quad \text{and} \quad \xi_0 + \frac{2}{2^{\nu_1}}$$

fall in the interval $(x_1; x_2)$; evidently, these numbers exist. Let

$$\xi_1 = \xi_0 + \frac{1}{2^{\nu_1}},$$

then since $\xi_1 = \frac{p}{q}$, where p is odd and $q = 2^{\nu_1}$ is divisible by 4, both series

$$\sum_{n=1}^{\infty} \frac{\sin^n \pi \xi_1 n}{n^\alpha}, \quad \sum_{n=1}^{\infty} \frac{\cos^n \pi \xi_1 n}{n^\alpha}$$

diverge to $+\infty$ by virtue of Theorems 1 and 2. We recurrently define $\nu_2, \nu_3, \dots$ so that

$$\nu_{k+1} = 2^{\nu_k+2}(A_{2^{\nu_k}} + k + 4) + 2\nu_k + 3,$$

where $A_{2^{\nu_k}}$ is provided by Lemma 5.1, and set

$$\xi_k = \xi_0 + \sum_{i=1}^k \frac{1}{2^{\nu_i}} \quad \text{and} \quad \xi = \xi_0 + \sum_{i=1}^{\infty} \frac{1}{2^{\nu_i}},$$

therefore $\xi \in (x_1; x_2)$, which satisfies our needs.

Now we are ready to show that for every positive integer k there exists such a number N_k that

$$\left| \sum_{n=1}^{N_k} \frac{\sin^n \pi \xi n}{n^\alpha} \right| > k \quad \text{and} \quad \left| \sum_{n=1}^{N_k} \frac{\cos^n \pi \xi n}{n^\alpha} \right| > k,$$

which means the desired divergence. We have

$$\begin{aligned} \left| \sum_{n=1}^{N_k} \frac{\sin^n \pi \xi n}{n^\alpha} \right| &\geq \left| \sum_{n=1}^{N_k} \frac{\sin^n \pi \xi_k n}{n^\alpha} - \sum_{n=1}^{N_k} \frac{\sin^n \pi \xi_k n - \sin^n \pi \xi n}{n^\alpha} \right| \geq \\ &\geq \left| \sum_{n=1}^{N_k} \frac{\sin^n \pi \xi_k n}{n^\alpha} \right| - \sum_{n=1}^{N_k} \frac{|\sin^n \pi \xi_k n - \sin^n \pi \xi n|}{n^\alpha}. \end{aligned}$$

Applying Lemma 5.1 with $q = 2^{\nu_k}$, $L = 2^{2q(A_q+k+4)}$, and $N = N_k = 2qL$ yields an estimate

$$\left| \sum_{n=1}^{N_k} \frac{\sin^n \pi \xi_k n}{n^\alpha} \right| \geq \frac{1}{2q} \log_2 L - A_q = k + 4$$

and the same one for the cos-series. Next, by the mean value theorem

$$\sum_{n=1}^{N_k} \frac{|\sin^n \pi \xi_k n - \sin^n \pi \xi n|}{n^\alpha} \leq \sum_{n=1}^{N_k} \frac{\pi n |\xi_k - \xi|}{n^\alpha} \leq \pi |\xi - \xi_k| \sum_{n=1}^{N_k} n \leq \pi N_k^2 |\xi - \xi_k|.$$

Since by construction

$$|\xi - \xi_k| \leq \frac{2}{2^{\nu_{k+1}}},$$

in line with the choice of ν_{k+1} we have

$$N_k^2 |\xi - \xi_k| \leq (2 \cdot 2^{\nu_k} \cdot 2^{2 \cdot 2^{\nu_k}(A_{2^{\nu_k}} + k + 4)})^2 \frac{2}{2^{2^{\nu_k+2}(A_{2^{\nu_k}} + k + 4) + 2\nu_k + 3}} = 1.$$

Thus

$$\left| \sum_{n=1}^{N_k} \frac{\sin^n \pi \xi n}{n^\alpha} \right| \geq k + 4 - \pi > k,$$

which is the required result. The same lower bound for the partial sum of the cos-series can be easily obtained quite the same way. $\square$

In conclusion we note that the set E from Theorem 5 consists solely of Liouville numbers.

References

- [1] Arkhipov G.I., Oskolkov K.I.: On a special trigonometric series and its applications. *Mathematics of the USSR. Sbornik*, **62:1**, 1989. 145–155.
- [2] Cassels J.W.S. An introduction to Diophantine approximation. *Cambridge Tracts in Mathematics and Mathematical Physics*, no. 45. Cambridge Univ. Press, New York, 1957.
- [3] Gelfond A. O.: Residues and their applications, Second edition, KomKniga, Moscow, 2006 (in Russian).
- [4] Hardy G.H., Littlewood J.E. Some problems of Diophantine approximation II. *Acta Mathematica*, **37**, 1914. 155–191.
- [5] Khinchin A. Ya. Continued Fractions. *Dover Publications*, Inc., New York, 1997.
- [6] Kolmogorov, A.N. Fomin, S.V. Introductory Real Analysis. *Dover Publications*, Inc., New York, 1975.
- [7] Laeng, E., Pata, V.: A convergence–divergence test for series of nonnegative terms. *Expositiones Mathematicae*, **29** (4), 2011. 420–424.
- [8] Salikhov, V.Kh., On the measure of irrationality of the number π , *Mathematical Notes*, **88:4**, 2010. 563–573.

Department of Mathematics and Mechanics, Moscow State University
1 Leninskie Gory, 119991, Moscow, Russia
`ab@rector.msu.ru`, `goryashin@mech.math.msu.su`